



Physical/Occupational Therapy Prescription

Name:

Date of Birth:

Diagnosis: Biceps Tendonitis

Code: M75.20

Procedure: Arthroscopic Biceps Tenodesis

Surgery Date:

Instructions: PT to begin after _____

Range of motion:

- Sling for the first 4 weeks post-operatively and then discontinue.
- Begin immediate range of motion including scapular range of motion.
- Progress from passive to active-assisted to active motion as tolerated.
- Begin with pendulums, pulleys, and wand/cane exercises.
- Avoid cross-body adduction and rotational motions in flexion or abduction until 140° elevation and 40° external rotation in adduction have been achieved.

Strengthening:

- Begin strengthening once pain has subsided and the patient is progressing towards symmetric active range of motion, which usually occurs at four weeks post-operatively.
- Begin with isometrics with the arm at the side and then progress to bands and light weights as tolerated. - Focus strengthening upon the rotator cuff, deltoid, and scapular stabilizers.
- Do not strengthen the rotator cuff more frequently than three times per week to avoid tendonitis. - Avoid positions of impingement during strengthening.
- Begin eccentrics, plyometrics, and sport-specific exercises at two months post-operatively.
- Return to athletics, including pitching, at three months.
- Begin throwing from the mound and collision sports at 4.5 months post-operatively

Limitations: No resisted elbow flexion or forearm supination for the first six weeks post-operatively to avoid stressing the biceps tenodesis.

Please provide a home exercise program

Modalities: Heat before and ice after therapy.

Frequency: 2-3 times/week

Duration: 6 weeks

Signature:

Date: